

Pakistan Must Act Now or Put Millions of Lives on a Risk of Antimicrobial Resistance

Faisal Rasheed

Life Science Group, Isotope Application Division, Pakistan Institute of Nuclear Science and Technology, Islamabad.

The review on antimicrobial resistance (AMR) prepared by a panel chaired by Jim O'Neill estimates that by 2050, 10 million human lives a year and 100 trillion US dollars of economy are at risk because of resistant infections and the impact of AMR will fall on low and middle-income countries.¹ The AMR has been detected in all parts of the world and is one of the serious threat and greatest challenge to global public health. World Health Organization (WHO) in 2014 presented data from 114 countries on AMR and documented in every region of the world.²

Pakistan has a high frequency of AMR in various commonly found pathogens including *Staphylococcus aureus*,³ *Acinetobacter baumannii*,⁴ *Pseudomonas aeruginosa*,⁵ *Escherichia coli*,⁶ *Salmonella typhi*,⁷ *Mycobacterium tuberculosis*,⁸ and *Helicobacter pylori*.⁹ The AMR is also reported by the authors from our neighboring countries including China,¹⁰ India¹¹ and Iran.¹²

There are number of causes associated with AMR specifically toward the antibiotics in Pakistan including lack of AMR surveillance at national level, quack doctors do not know and follow the recommended eradication therapies. The antibiotics are prescribed without performing susceptibility testing, self medication. There is a indiscriminate use of antibiotics by patients to treat the various infections and also un-necessary antibiotics use in livestock and agriculture.

The WHO conducted a survey in 2013-14 and reported that Pakistan has no national action plan to tackle AMR.¹³ Following the strategic objectives of the WHO Global Action Plan for AMR, the Ministry of National Health Services, Regulations and Coordination, Pakistan launched National Strategic Framework that is the first step to combat AMR.¹⁴ If State of Pakistan fails to tackle the AMR issue now, there will be more

devastating loss in terms of people and economy to this country in future. Therefore, the higher authorities and concern institutions should pay serious attention and huge commitment for combating AMR throughout the country using National Strategic Framework.

Conflict of interest: None declared.

References

1. Review on Antimicrobial Resistance. (Accessed on 11th June 2017) Available from URL:<http://amr-review.org/05.2016>
2. World Health Organization. Antimicrobial resistance: Global report on surveillance 2014. (Accessed on 15th December 2017) Available from URL: http://apps.who.int/iris/bitstream/10665/112642/1/9789241564748_eng.pdf/12.01.2015
3. Ullah A, Qasim M, Rahman H, Khan J, Haroon M, Muhammad N, et al. High frequency of methicillin-resistant *Staphylococcus aureus* in Peshawar Region of Pakistan. Springerplus2016; 5: 600.
4. Hasan B, Perveen K, Olsen B, Zahra R. Emergence of carbapenem-resistant *Acinetobacterbaumannii* in hospitals in Pakistan. J Med Microbiol 2014; 63: 50-5.
5. UllahW, Qasim M, Rahman H, Bari F, Khan S, Rehman ZU, et al. Multi drug resistant *Pseudomonas aeruginosa*: Pathogen burden and associated antibiogram in a tertiary care hospital of Pakistan. Microb. Pathog 2016; 16: 209-12.
6. Rana F, Siddiqui S, Khan A, Siddiqui F, Noreen Z, Bokhari Set al. Resistance patterns of diversified phylogroups of *Escherichia coli* associated with mothers having history of preterm births in Pakistan. J Matern Fetal Neonat Med 2017; 30: 68-73.
7. Hasan R, Zafar A, Abbas Z, Mahraj V, Malik F, Zaidi A. Antibiotic resistance among *Salmonella enteric serovarstyphi* and *Paratyphi A* in Pakistan (2001-2006). J Infect Dev Ctries 2008; 2(4): 289-94.
8. Ali A, Hasan Z, McNerney R, Mallard K, Hill-Cawthorne G, Coll F, et al. Whole genome sequencing based characterization of extensively drug-resistant *Mycobacterium tuberculosis* isolates from Pakistan. PLoS One 2015; 10:e0117771. doi:10.1371/journal.pone.0117771.
9. Rasheed F, Campbell BJ, Alfizah H, Varro A, Zahra R, Yamaoka Y, et al. Analysis of clinical isolates

Corresponding Author:

Faisal Rasheed

Life Science Group, Isotope Application Division
Pakistan Institute of Nuclear Science and Technology, Islamabad.
Email: frohpl@gmail.com

Received: 13 December 2016, Accepted: 07 December 2017,
Published: 22 December 2017

- of *Helicobacter pylori* in Pakistan reveals high degrees of pathogenicity and high frequencies of antibiotic resistance. *Helicobacter* 2014; 19: 387–99.
- 10. Jianyong Wu, Lanlan Liu, Wang G Jiahai Lu. One Health in China. *Infect. Ecol. Epidemiol* 2016; 6: 10.
 - 11. Kumar SG, Adithan C, Harish BN, Sujatha S, Roy G, Malini A. Antimicrobial resistance in India: A review. *J Nat Sci Biol Med* 2013; 4(2): 286-91.
 - 12. Mahmoudi S, Pourakbari B, Moradzadeh M, Eshaghi H, Ramezani A, Haghi AMT, et al. Prevalence and antimicrobial susceptibility of *Salmonella* and *Shigella* spp. among children with gastroenteritis in an Iranian referral hospital. *MicrobPathog* 2017; 109: 45-8.
 - 13. World Health Organization. Antimicrobial resistance. (Accessed on 11th June 2017) Available from URL:http://who.int/drugresistance/documents/situation_analysis/29.04.2015
 - 14. NHSRC. National Strategic Framework for Anti-Microbial Resistance. (Accessed on 11th June 2017) Available from URL: http://www.nhsrc.gov.pk/news_details977f.html?news_id=228
-